

SHORT COMMUNICATION

Phytotoxicity, cytotoxicity and antioxidant activity of the invasive shrub *Austro eupatorium inulifolium* (Kunth) R.M. King & H. Rob

Indika Chandrasiri¹, Shashini Diwakara¹, Chamara Janaka Bandara¹, Siril Wijesundara², Sumedha Madawala^{3*} and Veranja Karunaratne^{1,4}

¹Department of Chemistry, University of Peradeniya, Peradeniya, Sri Lanka.

²Royal Botanic Gardens, Peradeniya, Sri Lanka.

³Department of Botany, University of Peradeniya, Peradeniya, Sri Lanka.

⁴Sri Lanka Institute of Nanotechnology, Homagama, Sri Lanka.

Accepted December 31, 2015

ABSTRACT

Methanol and dichloromethane extracts of root, stem, leaves and flowers of invasive plant *Austro eupatorium inulifolium* were tested for cytotoxic, phytotoxic, antioxidant and antifungal activities. Significantly higher phytotoxicity was detected in methanol extracts of leaves and dichloromethane extracts of roots. This effect was most pronounced against amaranth seeds where the seed germination was reduced to 3.74 % with the addition of methanol extracts of leaves at 3000 mg/L. The dichloromethane extract of roots of *A. inulifolium* showed potent antifungal activity against *Cladosporium cladosporioides*. Cytotoxic activity was found in dichloromethane extract of roots ($LD_{50} = 27.91 \pm 8.55$ mg/L), methanol extract of flowers ($LD_{50} = 15.22 \pm 7.89$ mg/L) and leaves ($LD_{50} = 22.92 \pm 11.76$ mg/L) against *Artemiasalina* (brine shrimp) larvae. The results also revealed significant antioxidant activity in methanol extract of leaves ($IC_{50} = 33.66 \pm 0.03$ mg/L) against the reference α -tocopherol ($IC_{50} = 10.02 \pm 0.01$ mg/L).

Keywords: invasive plant, cytotoxicity, phytotoxicity, antioxidant and antifungal activities.

INTRODUCTION

Plants have co-evolved with their respective environments for millions of years. During this process, secondary metabolites played a critical role in plants' adaptations to insect and microbial attacks and climatic conditions. Plant metabolites with various bioactivities including phytotoxic and antioxidant activities allow plants to live in a state of equilibrium with the environment. However, some plant species use these bioactivities to transform themselves into noxious weeds. The indigenous flora of Sri Lanka comprises about 7,500 species. Of the 3,154 flowering plants, about 894 (28 %) species are endemic to the island, some carrying varying levels of biological activity (Wijesundara *et al.*, 2012; Bandara *et al.*, 1989a; Bandara *et al.*, 1990a; Hewage *et al.*, 1997; Hewage *et al.*, 1998). Many bioactivities in plants have been used in medicinal applications (Banadara *et al.*, 1989b; Bandara *et al.*, 1989c; Bandara *et al.*, 1990b; Williams *et al.*, 2011). Plant ecologists were always puzzled by the fact that exotic plant species with low densities in their native ranges eventually produce large populations in their introduced ranges. The 'novel weapons hypothesis' proposed that these exotic plants possess novel biochemical compounds with powerful bioactivities (Callaway and Ridenour, 2004). *Austro eupatorium inulifolium* (Kunth)

King and Robinson (Asteraceae) which is an aggressive invasive shrub native to South America has become a noxious invader in the up-country, wet zone of Sri Lanka, invading many natural and man-made ecosystems (Madawala *et al.*, 2014). It is also listed as an 'agricultural and environmental weed in the Global Compendium of weeds (Randall, 2012). Even though it has been well identified as an invasive plant in many countries, it has been categorized as an invasive plant in Sri Lanka fairly recently. *A. inulifolium* has been invading the *Cymbopogon*-dominated grasslands in the Knuckles Conservation Area (KCA), displacing the grass. Once established, *A. inulifolium* can form mono-specific stands in their introduced range, influencing the growth and survival of the native flora (Haluwana and Madawala, 2013).

There have been many hypotheses to explain the mechanisms of exotic plant invasions into new landscapes and among them the 'novel weapon hypothesis' has attracted contrasting views since it was proposed in 2004 (Blair *et al.*, 2006; Duke *et al.*, 2009). According to this hypothesis, some exotic species credit their success of spread due to the production of bioactive compounds that native species never encountered before (Callaway and Ridenour, 2004; Thorpe *et al.*, 2009; Callaway and Aschehoug, 2000). Previous studies on other

*Corresponding author's email: sumedha.madawala@gmail.com

invasive species have confirmed the presence of bioactive compounds (Yan *et al.*, 2010; Shao *et al.*, 2010; Ens *et al.*, 2009; Xie *et al.*, 2010). These bioactive compounds can reach the soil environment through leaching, root exudation or litter decomposition. In the present study, we attempted to investigate the potential role of bioactive compounds in *A. inulifolium* by using bioassays with extracts of different parts of the shrub including stems, roots, flowers and leaves and to relate them to its invasive success. Apart from the isolation of nine new norlabdane derivatives with antimicrobial activity in *A. inulifolium* (Saito *et al.*, 2011), no studies have been carried out to assess the bioactive profile of *A. inulifolium* with an aim to explain its invasive nature of the plant. We report herein, the bioactivities of *A. inulifolium* including cytotoxic, phytotoxic, antioxidant and antifungal properties.

MATERIALS AND METHODS

Austroeupeatorium inulifolium, belongs to the family Asteraceae, is a shrub that can grow up to 1-5 m in height. It bears a creamy white fragrant inflorescence (Figure 1). Leaves are opposite, spear-shaped and pubescent, abruptly narrowing to a wedge-shaped petiole. It is well known as an aggressive invader in many countries.



Figure 1. The creamy white inflorescence of *A. inulifolium*. (Photo Credit: InokaPiyasinghe).

Extraction of potential bioactive compounds

Plant samples were collected from Riverston area in the Knuckles Forest Reserve (KFR) in the central of Sri Lanka. Plant parts (roots, stems, leaves and flowers) were separated and air-dried before grinding into a powder. The resulting powdered plant material (≈ 600 g) was subjected to

sequential extraction with 5 L of CH_2Cl_2 and CH_3OH separately after maceration for 24 hours (in cleaned and dried glass bottles) with constant shaking. The extract was then filtered and concentrated using a rotary evaporator at 30°C to obtain the final yield of extract in paste form. The crude extract was then stored at 4°C until further use. The CH_2Cl_2 and CH_3OH crude extracts were subjected to cytotoxic, phytotoxic, antioxidant and antifungal bioassays.

Cytotoxic Activity

Cytotoxic activity was determined using a brine shrimp lethality assay (McLaughlin, 1982). Brine shrimp eggs were added to fresh artificial sea water and allowed to hatch in a beaker which was aerated and illuminated using a 20 W bulb. The temperature was maintained at 27°C . A series of concentrations (0.75, 2, 7.5, 20, 75, 200, 750, 2000 mg/L) of the dichloromethane and methanol extracts of different plant parts were tested with 1-day old brine shrimps (three replicates and 10 shrimps per experiment). After 24 hours, the number of surviving nauplii of brine shrimps was counted and LD_{50} value (concentration at 50 % survival) was calculated using probit analysis of MINITAB version 16. The known cytotoxic lactone, (4S)-4-methyl-2-(11-dodecynyl)-2-butenolide, from the genus *Hortonia* was used as the positive control (Ratnayake *et al.*, 2001).

Phytotoxic activity

The phytotoxic activity was determined using a seed assay with two monocot species, maize (*Zea mays*) and bluegrass (*Poa sp.*), and two dicot species, amaranth (*Amaranthus sp.*) and radish (*Raphanus sativus*). Ten seeds (surface sterilized with 0.5 % HgCl_2) from each species were treated with 5 ml of 250, 750 and 3000 mg/L of the CH_2Cl_2 and CH_3OH extracts of different plant parts in petri dishes (9 cm diameter) lined with Whatman No. 3 filter paper. Distilled water mixed with 1 % DMSO (v/v %) was used as the negative control and [(4S)-4-methyl-2-(11-dodecynyl)-2-butenolide] as the positive control. Petri dishes were sealed with Parafilm to prevent any water losses and the petri plates were stored in the dark at room temperature. Tests were carried out in triplicate for each concentration. Cumulative germination was determined by counting the number of germinated seeds daily over a period of 6 days and the Germination Index was calculated using the formula (Chiapuso *et al.*, 1997),

$$\text{GI} = (N_1) \times 1 + (N_2 - N_1) \times 1/2 + (N_3 - N_2) \times 1/3 + \dots + (N_n - N_{n-1}) \times 1/n,$$

where $N_1, N_2, N_3 \dots N_n$: proportion of germinated seeds observed afterwards 1, 2, 3... n-1 and n is the number of days. On the 7th day, shoot and root lengths of the seedlings were measured. The inhibitory or stimulatory percentage was calculated using the

equation (Chung *et al.*, 2001), Inhibition (-) or Stimulation (+) % = (Extract – Control)/(Control) × 100.

Antioxidant activity

Antioxidant activity was determined using 2, 2-diphenyl-1-picrylhydrazyl (DPPH) radical scavenging assay (Budzianowski and Budzianowska, 2006). The reduction of DPPH in the extract was measured using a concentration series ranging from 1 – 500 mg/L at 517 nm using a UV-visible spectrophotometer (Shimadzu UV-1800) after 30 minutes against the CH₃OH blank. The initial absorbance of DPPH and each test solution (without DPPH) were measured at 517 nm separately. α -Tocopherol (Vitamin E), concentration series ranging from 1 – 500 mg/L was used as the positive control. Tests were carried out in triplicate for each concentration. Percentage antioxidant activity was calculated using absorbance readings using the equation, $[A_0 - A_t/A_t] \times 100$ where A_0 is the initial absorbance value and A_t is the absorbance of the test sample after 30 min of adding DPPH. The % antioxidant activity was plotted against the concentration gradient for all three trials separately and trend lines were constructed for the linear range of the plot. The IC₅₀ value was calculated for each trend line from which the average was taken.

Antifungal activity

Quantitative analysis of antifungal activity using the TLC bioassay method with the fungus, *Cladosporium cladosporioides* was carried out. A concentration series of the extracts (250, 750 and 2500 mg/L) of different plant parts was prepared and spotted on a TLC plate. A conidial suspension of *C. cladosporioides* was carefully sprayed using an atomizer onto the TLC plates. After 2 days of incubation, the presence of antifungal compounds was detected using inhibition zones. Antifungal active spots appeared white against the background of grey-green mycelia.

RESULTS AND DISCUSSION

Yields of potential bioactive extracts

After extraction and concentration, crude yields of methanol and dichloromethane extracts of different plant parts of *A. inulifolium* were obtained (Table 1). In both extracts, the highest percentage yield was obtained from the leaves.

Cytotoxic effects

Moderate cytotoxic activity was detected in the CH₂Cl₂ extract of roots (27.91 ± 8.55 mg/L), and CH₃OH extracts of leaves (22.92 ± 11.76) and flowers (15.22 ± 7.89). The LD₅₀ value of the positive control, (4S)-4-methyl-2-(11-dodecynyl)-2-butenolide, was 0.80 mg/L (± 0.37).

Phytotoxic activity

Germination index (GI) generally decreased with increasing concentrations of CH₃OH and CH₂Cl₂ extracts of *A. inulifolium* (Table 2). However, CH₃OH-leaf, CH₂Cl₂-root and CH₂Cl₂-stem extracts showed the most significant reductions in germination index with increasing concentrations (from 250 to 3000 mg/L). As an example, CH₃OH-leaf extracts decreased the GI from about 68 % to 4 % in amaranth seeds, while CH₂Cl₂-root extracts reduced the GI of same seeds from 61 % to 8 %, when increase the concentration from 250 mg/L to 3000 mg/L. In contrast, all concentrations of CH₂Cl₂-flower extracts showed no influence on the germination of seeds tested in the assay. CH₂Cl₂-leaf extracts too showed no impact on the germination of maize seeds.

Both CH₃OH and CH₂Cl₂ extracts of *A. inulifolium* roots and shoots negatively influenced the growth of both monocot and dicot seedlings tested, and this inhibition gradually increased with increasing concentrations (Figure 2). CH₃OH extracts of *A. inulifolium* leaves negatively affect the growth of seedlings while CH₂Cl₂ extracts did not show clear inhibition apart from *amaranth* seedlings.

Table 1. Mass and the yield (as a percentage) of methanol and dichloromethane crude extracts of different plant parts of *A. inulifolium* collected from Riverston area in the Knuckles Forest Reserve, Sri Lanka.

Plant part	Dry mass (g)	Mass of CH ₃ OH crude extract (± 0.001 g)	% yield of CH ₃ OH crude extract	Mass of CH ₂ Cl ₂ crude extract (± 0.001 g)	% yield of CH ₂ Cl ₂ crude extract
Flower	77	4.14	5.37	0.67	0.86
Leaf	283	32.07	11.33	20.49	7.24
Stem	410	4.65	1.13	1.02	0.25
Root	173	3.41	1.97	2.15	1.24

Table 2. Germination Index (GI, %) of monocot (maize and bluegrass) and dicot (amaranth and radish) seeds under different concentrations (250, 750 and 3000 mg/L) of plant extracts of *A. inulifolium*.

Extract	Conc. (mg/L)	Monocots		Dicots	
		Maize	Bluegrass	Amaranth	Radish
CH ₂ Cl ₂ -leaf	250	98.1 ^a	88.1 ^a	83.6 ^a	95.0 ^a
	750	90.6 ^a	77.2 ^{ab}	71.7 ^b	86.2 ^b
	3000	89.9 ^a	69.4 ^b	43.2 ^c	71.5 ^c
CH ₃ OH-leaf	250	98.9 ^a	89.5 ^a	67.6 ^a	79.4 ^a
	750	91.3 ^a	68.8 ^b	46.8 ^b	68.3 ^b
	3000	74.8 ^b	60.8 ^b	3.7 ^c	44.1 ^c
CH ₂ Cl ₂ -flower	250	101.8 ^a	98.7 ^a	92.5 ^a	92.2 ^a
	750	97.0 ^a	93.3 ^a	90.3 ^a	88.7 ^a
	3000	88.6 ^a	91.9 ^a	89.5 ^a	86.1 ^a
CH ₃ OH-flower	250	95.1 ^a	92.7 ^a	89.1 ^a	90.0 ^a
	750	87.6 ^{ab}	88.3 ^a	72.5 ^b	72.1 ^b
	3000	80.9 ^b	77.4 ^b	48.3 ^c	66.8 ^c
CH ₂ Cl ₂ -root	250	99.9 ^a	87.4 ^a	61.4 ^b	98.3 ^a
	750	69.9 ^b	57.8 ^b	42.0 ^c	91.7 ^a
	3000	61.1 ^b	35.3 ^c	8.2 ^d	75.0 ^b
CH ₃ OH-root	250	89.2 ^{ab}	61.5 ^b	68.3 ^b	100.0 ^a
	750	63.8 ^c	63.3 ^b	58.0 ^{bc}	92.8 ^b
	3000	74.1 ^{bc}	30.5 ^c	51.1 ^c	93.9 ^b
CH ₂ Cl ₂ -stem	250	80.3 ^{ab}	72.8 ^b	98.3 ^a	61.3 ^b
	750	65.2 ^b	52.4 ^c	52.9 ^c	89.2 ^{ab}
	3000	40.9 ^c	8.8 ^d	7.8 ^d	76.4 ^b
CH ₃ OH-stem	250	90.2 ^{ab}	75.3 ^b	58.2 ^b	98.3 ^{ab}
	750	85.1 ^b	59.9 ^b	56.3 ^b	92.2 ^{ab}
	3000	49.7 ^c	25.9 ^c	35.0 ^c	92.8 ^b

Means with the same letters in a column (between different concentrations) indicate no significant differences at $P < 0.05$.

Interestingly, CH₃OH-leaf and CH₃OH-flower extracts showed higher inhibition on the growth of dicot seedlings compared to that of monocots. In contrast to the effects of other extracts, CH₂Cl₂-flower extracts stimulated the growth of dicot and monocot seedlings, showing highest stimulatory effect at 250 mg/L concentration, though the differences are not significant. Furthermore, the monocot seedlings were more stimulated than that of dicot seedlings (Figure 2).

Antioxidant activity of *A. inulifolium*

When comparing the antioxidant activity of

CH₃OH and CH₂Cl₂ extracts of *A. inulifolium* with α -tocopherol ($IC_{50} = 10.02 \pm 0.01$ mg/L), the CH₃OH extract of leaves ($IC_{50} = 33.66 \pm 0.03$ mg/L) showed a significant antioxidant activity among the extracts at $P < 0.05$.

Antifungal activity of *A. inulifolium*

Interestingly, the antifungal activity was only observed in CH₂Cl₂-root extracts. According to the results obtained, the average diameters of inhibition zones shown in the presence of the CH₂Cl₂ extract of *Austroeupeatorium* roots were 16.0 and 24.6 mm at concentrations 750 and 2500

mg/L, respectively (Table 3).




Figure 2. Inhibitory or Stimulatory effects (%) of different concentrations (250, 750 and 3000 mg/L) of *A. inulifolium* extracts on the growth of monocot (maize and bluegrass) and dicot (amaranth and radish) seedlings.

Different letters indicate significant differences among concentrations at $P < 0.05$.




Figure 3. Inhibition zones given by CH_2Cl_2 extract of *A. inulifolium* roots (a) 750 mg/L (b) 2500 mg/L.

Table 3. Diameters of the inhibition zones given by different concentrations of root dichloromethane extract of *A. inulifolium* (2500 mg/L and 750 mg/L).

Concentration/(mg/L)	Diameter of the inhibition zone (± 0.5 /mm)		
	Trial 1	Trial 2	Trial 3
2500	24.0	27.0	23.0
750	16.0	14.0	18.0

DISCUSSION

When consider the extraction of potentials of bioactive compounds in *A. inulifolium*, the CH_3OH extracts recorded a higher yield than that of CH_2Cl_2 extracts. This may be due to the higher content of polar compounds in *A. inulifolium* compared to that of non-polar compounds. Cytotoxic results suggested that *A. inulifolium* roots, flowers and leaves contain cytotoxic compounds which can be useful as antiproliferative and antitumor activities, as well as pesticidal and other bioactive agents. There is evidence to support that natural cytotoxic compounds are good contenders for anticancer drugs (Rahman *et al.*, 2001). In addition, these cytotoxic compounds also play an important role in these plants to minimize pest attacks compared to co-existing natives, determining their successful establishment in their introduced ranges (McGraw and Eloff, 2008).

The effects of bioactive compounds on seed germination have been tested using germination index (GI) in previous studies (Chiapuso *et al.*, 1997). GI is considered as a sensitive indicator of phytotoxic effects (Ahmed and Wardle, 1994). Results showed that germination of tested species were affected by all extracts which either delayed

or reduced germination of all test seeds. However, a significantly higher phytotoxic effect was detected in CH_3OH -leaf and CH_2Cl_2 -root extracts of *A. inulifolium* ($P < 0.001$). The CH_3OH -leaf extract at 3000 mg/L showed the most pronounced impact against amaranth seeds with GI of 3.74 %. The analysis showed that CH_2Cl_2 -flower extract had no significant effect on the GI ($P < 0.05$) of both monocot and dicots, although it has declined the seed germination. The degree of effect on germination increased with the increasing concentrations of extracts and this was evident in extracts of all plant parts of *A. inulifolium*. The delayed seed germination can have important biological implications, as this can negatively affect the establishment of seedlings of co-occurring natives (Chaves *et al.*, 2001), and thereby increasing the chances of competing for resources with neighbouring species (Xingxinag *et al.*, 2009).

However, the effect was more pronounced in CH_3OH extracts of *A. inulifolium*, where polar compounds are commonly found. Methanolic extracts of some plant species were known to contain phenolics and other toxic substances (Belicova *et al.*, 2001; Al Harun *et al.*, 2015). These phenolics may inhibit the germination process through their interference in the indole acetic acid metabolism, or synthesis of protein and

iron uptake by plants (Blum, 1998; Castro *et al.*, 1984).

Phytotoxic results showed that the root and shoot development of germinating test seeds are highly inhibited by the CH₂Cl₂ extracts of root and stem, and by all extracts of CH₃OH. In contrast, slight stimulatory effects on the root and shoot growth of monocot seedlings was shown by CH₂Cl₂ extracts of leaves and flowers. The shoot growth appeared less sensitive to *A. inulifolium* extracts than the radicle, which eventually grows into roots. The phytotoxic compounds are highly active against meristematic tissue in growing roots. This perhaps is the reason for differing responses of root and shoot to *A. inulifolium* extracts. Phytotoxicity has been suggested as the key strategy for the impressive success of many invasive plants that has been dominated in their invaded plant communities (Ridenour and Callaway, 2001).

In some instances, there were slight stimulatory effects on root and/or shoot by different extracts of *A. inulifolium*. According to results, inhibition or stimulation of root and shoot and the germination index varied with plant extract, concentration and seed type. The stimulatory bioactive compounds can be used to develop eco-friendly, cheap and effective Green Growth Promoters (Oudhia *et al.*, 1998). Phytotoxic compounds inhibit germination and seedling growth probably by affecting cell division and elongation, processes that are very important at early stages, or by interfering with enzymes involved in the mobilization of nutrients necessary for germination (Batlang and Shush, 2007).

Antioxidant results revealed that the MeOH extract of leaves contains compounds with antioxidant activity. Generally, plant polyphenolic compounds are responsible for antioxidant activity. In addition, they also act as iron chelators. The latter property may be important in an invasive plant like *A. inulifolium* where it contributes to altering the soil bacterial flora in their growing habitats (Morel *et al.*, 1993).

In the antifungal assay, larger inhibition zones were observed with CH₂Cl₂ extracts, suggesting that the solvent has the potential to extract many active antifungal compounds. Chromene, an antifungal compound, found in roots of the invasive plant *Eupatorium riparium* which also belongs to the same genus as *A. inulifolium* (Bandara *et al.*, 1992). Furthermore, the antifungal, phytotoxic and antioxidant activities were also tested in essential oils extracted from roots and inflorescence of *Eupatorium adenophorum* (Ahluwalia *et al.*, 2014), further supporting this genus's worldwide distribution.

CONCLUSIONS

The results suggest that *A. inulifolium* contained phytotoxic, cytotoxic and antifungal activities. The presence of phytochemicals in *A. inulifolium* may suppress the growth of other plants in the habitat they invade. Cytotoxic compounds can kill pests thereby minimizing pest attacks. Antifungal compounds can reduce fungal attacks on the plant too. These characteristics can favour invasive species to invade and establish in their introduced ranges successfully. Therefore, the survival and spread of *A. inulifolium* may have been favored by the presence of these phytotoxic compounds where they can inhibit the growth of co-habiting plants. These factors may have contributed to the aggressive nature of *A. inulifolium*.

REFERENCES

- Ahluwalia, V., Sisodia, R., Walia, S., Sati, O.P., Kumar, J. and Kumdu, A. (2014). Chemical analysis of essential oils of *Eupatorium adenophorum* and their antimicrobial, antioxidant and phytotoxic properties. *Journal of Pest Science* **87** (2): 341-349.
- Ahmed, M. and Wardle, D.A. (1994). Allelopathic potential of vegetative and flowering ragwort (*Senecio jacobaea* L.) plants against associated pasture species. *Plant and Soil* **164**(1): 61-68.
- Bandara, B.M.R., Fernando, I.S.H., Hewage, C.M., Karunaratne, V., Adikaram, N.K.B. and Wijesundara, D.S.A. (1989a). Antifungal activity of some medicinal plants of Sri Lanka. *Journal of the National Science Council of Sri Lanka* **17**(7): 1-13.
- Bandara, B.M.R., Hewage, C.M., Karunaratne, V. and Adikaram, N.K.B. (1989b). Methyl ester of *para*-coumaric acid: antifungal principle of the rhizome of *Costusspeciosus*. *Planta Medica* **5**: 477-478.
- Bandara, B.M.R., Cortez, D., Jayasinghe, L., Karunaratne, V., Wannigama, G.P. and Sotheeswaran, S. (1989c). Aphorphine alkaloids from *Litsea gardineri* and *Actinodaphne speciosa*. *Planta Medica* **55**: 393.
- Bandara, B.M.R., Hewage, C.M., Jayamanne, D.H.L.W., Karunaratne, V., Adikaram, N.K.B., Pinto, M.R.M. and Wijesundara, D.S.A. (1990a). Biological activity of some steam distillates from ten species of Rutaceae Plants. *Journal of the National Science Council of Sri Lanka* **18** (1): 71.
- Bandara, B.M.R., Jayasinghe, L., Karunaratne, V., Wannigama, G.P., Bokel, M., Kraus, W. and Sotheeswaran, S. (1990b). Isolation of pheanthine from *Cyclea burmanai*. *Planta Medica* **56**: 245.

- Bandara, B. M. R., Hewage, C. M., Karunaratne, V., Wannigama, G. P. and Adikaram, N. K. B. (1992). An Antifungal Chromene from *Eupatorium riparium*. *Phytochemistry* **31**: 1993.
- Batlang, U. and Shushu, D.D. (2007). Allelopathic activity of sunflower (*Helianthus annuus* L.) on growth and nodulation of bambara groundnut (*Vigna subterranean* (L.) Verdc.). *Journal of Agronomy* **6**(4): 541-547.
- Belicova, A., Krizkova, L., Nagy, M., Krajcovic, J. and Ebringer, L. (2001). Phenolic acids reduce the genotoxicity of acridine orange and ofloxacin in *Salmonella typhurium*. *Folia Microbiologica* **46**(6): 511-514.
- Blair, A.C., Nissen, S.J., Brunk, G.R. and Hufbauer, R.A. (2006). A lack of evidence for an ecological role of the putative allelochemical (\pm)-catechin in spotted knapweed invasion success. *Journal of Chemical Ecology* **32**:2327–2331.
- Blum, U. (1998). Effects of microbial utilization of phenolic acids and their phenolic acid breakdown products on allelopathic interactions. *Journal of Chemical Ecology* **24**: 685-708.
- Budzianowski, J. and Budzianowska, A. (2006). Chromatographic and spectrophotometric analyses of the DPPH free radical scavenging activity of the fractionated extracts from *Lamium album* L., *Lamium purpureum* L. & *Viscum album* L. *Herba Polonica* **52**(1/2): 51-57.
- Callaway, R.M. and Aschehoug, E.T. (2000). Invasive plants versus their new and old neighbors: A mechanism for exotic invasion. *Science* **290**:521–523.
- Callaway, R.M. and Ridenour W.M. (2004). Novel weapons: a biochemically based hypothesis for invasive success and the evolution of increased competitive. *Frontiers in Ecology and the Environment* **2**:436–433.
- Castro, P.R.C., Rodrigues, J.C., Viegas, R.F.A., Lima, G.P.P., Jureidin, P. and Denbandal, I.M. (1984). Allelopathic action of some weed extracts on rice (*Oryza sativa* L.). *Agriculturist* **41**(1): 369-381.
- Chaves, N., Sosa, T. and Escudero, J.C. (2001). Plant growth inhibiting flavonoids in exudate of *Cistus ladanifer* and in associated soils. *Journal of Chemical Ecology* **27** (3): 623-631.
- Chiapuso, G., Sanchez, A. M., Reigosa, M.J., Gonzalez, L. and Pellissier, F. (1997). Do germination indices adequately reflect allelochemical effects on the germination process? *Journal of Chemical Ecology* **23**(11): 2445- 2453.
- Chung, I.M., Ahn, J.K. and Yun, S.J. (2001). Assessment of allelopathic potential of barnyard grass (*Echinochloa crus-galli*) on rice (*Oryza sativa* L.) cultivars. *Crop Protection* **20**(10): 921-928.
- Duke, S.O., Blair, A.C., Dayan, F.E., Johnson, R.D., Meepagala, K.M., Cook, D. and Bajsa J. (2009). Is catechin a novel weapon of spotted knapweed (*Centaurea stoebe*)? *Journal of Chem Ecology* **35**:141–153.
- Ens, E.J., Bremner, J.B., French, K. and Korth, J. (2009). Identification of volatile compounds released by roots of an invasive plant, bitou bush (*Chrysanthemoides monilifera* spp. *rotundata*), and their inhibition of native seedling growth. *Biological Invasions* **11**:275–287.
- Al Harun M.A.Y., Johnson J., Uddin M.N. and Robinson R.W. (2015). Identification and Phytotoxicity Assessment of Phenolic Compounds in *Chrysanthemoides monilifera* subsp. *monilifera* (Boneseed). *PLoS ONE* **10**(10): e0139992. doi:10.1371/journal.pone.0139992.
- Haluwana, N. and Madawala, H.M.S.P. (2013). Changes in Plant Diversity and Composition across Forest Edges Bordered by *Austro eupatorium inulifolium* Invaded Grasslands in the Knuckles Conservation Area, Sri Lanka. *Ceylon Journal of Science (Biological Sciences)* **42**(2): 29-43.
- Hewage, C.M., Bandara, K.A.N.P., Karunaratne, V., Bandara, B.M.R. and Wijesundara, D.S.A. (1997). Insecticidal Activity of some Medicinal Plants of Sri Lanka. *Journal of the National Science Council of Sri Lanka* **25** (3):141 – 150.
- Hewage, C.M., Bandara, B.M.R., Karunaratne, V., Wannigama, G.P., Pinto, M.R.M. and Wijesundara, D.S.A. (1998). Antibacterial activity of some medicinal plants of Sri Lanka. *Journal of the National Science Foundation of Sri Lanka* **26**(1): 27-34.
- Kitsati, N., Fokas, D., Oyuzouni, M-D., Mantzaris, M.D., Barbouti, A. and Galaris, D. (2012). Lipophilic caffeic acid derivatives protect cells against H₂O₂-induced DNA damage by chelating intracellular iron. *Journal of Agricultural and Food Chemistry* **60**: 7873-07879.
- Madawala, S., Chandrasiri, I., Diwakara, S., Wijesundara, S. and Karunaratne, V. (2014). Bioactivities of invasive plant *Austro eupatorium Inulifolium*, *Proceedings of the University of Peradeniya Research Sessions* **18**: 477.
- McGraw, L.J. and Eloff, J. N. (2008). Ethnoveterinary use of Southern African plants and scientific evaluation of their medicinal properties. *Journal of Ethnopharmacology* **119**:559-574.
- McLaughlin, J.L. (1982). Brine shrimp: a convenient general bioassay for active

- constituents. *Planta Medica* **45**: 31-32.
- Morel, I., Lescoat, G., Cogrel, P., Sergent, O., Padeloup, N., Brissot, P., Cillard, P. and Cillard, J. (1993). Antioxidant and iron-chelating activities of the flavonoids catechin, quercetin and diosmetin on iron-loaded rat hepatocyte cultures. *Biochemical Pharmacology* **45**(1):13-9.
- Oudhia, P., Kolhe, S.S. and Tripathi, R.S. (1998). Allelopathic effect of *Blumealacera* L. on rice and common Kharif weeds. *Oryza* **35**(2), 175-177.
- Rahman, M.M., Habib, M.R., Hasan, S.M.R., Sayeed, M.A. and Rana, M.S. (2011). Antibacterial, cytotoxic and antioxidant potential of methanolic extract of *Phyllanthusacidus* L. *International Journal of Drug Development and Research* **3**(2): 154-161.
- Randall, R.P. (2012). A global compendium of weeds. 2nd Edition. Department of Agriculture and Food, Western Australia.
- Ratnayake, R., Karunaratne, V., Bandara, B. M. R., Kumar, V., Macleod, J.K. and Simmonds, P. (2001). Two new lactones with mosquito larvicidal activity from three *Hortonia* species. *Journal of Natural Products* **61**:376-378.
- Ridenour, W.M. and Callaway, R.M., (2001). The relative importance of allelopathy in interference: the effects of an invasive weed on a native bunchgrass. *Oecologia* **126**(3), 444-450
- Saito, Y., Matsuo, S., Sutoyo, A. and Tori, M. (2011). Nine new norlabdane diterpenoids from the leaves of *Austro eupatorium inulifolium*. *Helvetica Chimica Acta* **94** (2): 313 -326.
- Shao, H., Peng, S.L., Wei, X.Y., Zhang, D.Q. and Zhang, C. (2010). Potential bioactive compounds from an invasive weed *Mikaniamicrantha* H.B.K. *Journal of Chemical Ecology* **31**:1657-1668.
- Thorpe, A.S., Thelen, G.C., Diaconu, A. and Callaway, R.M. (2009). Root exudate is allelopathic in invaded community but not in native community: Field evidence for the novel weapons hypothesis. *Journal of Ecology* **97**:641-645.
- Wijesundara, S., Kathriarachchi, H.S., Ranasinghe, S.W. and Hapuarachchi, G. (2012). Analysis of seed plants of Sri Lanka. In: *The National Red List 2012 of Sri Lanka; Conservation Status of the Fauna and Flora*. Weerakoon, D.K. & S. Wijesundara Eds., Ministry of Environment, Colombo, Sri Lanka. 340-345 pp.
- Williams, D. E., Loganzo, F., Whitney, L., Togias, J., Harrison, R., Singh, M.P., McDonald, L.A., Kathirgamanathar, S., Karunaratne, V. and Andersen, R.J. (2011). Depsides isolated from the Sri Lankan lichen *Parmotrema* sp. exhibit selective Plk1 inhibitory activity. *Pharmaceutical Biology* **49**(3):296-301.
- Xie, L.J., Zeng, R.S., Bi H.H., Song, Y.Y., Wang, R.L., Su, Y.J., Chen, M., Chen, S. and Liu, Y.H. (2010). Allelochemical mediated invasion of exotic plants in China. *Allelopathy Journal* **25**:31-50.
- Xingxinag, G., Li Zongjung, M.G., Changsong, L. and Zuowen, S. (2009). Allelopathic effects of *Hemister palyrata* on the germination and growth of wheat, sorghum, cucumber, rape and radish seeds. *Weed Biology and Management*. **9**: 243 - 249.
- Yan, J., Bi, H.H., Liu, Y.Z., Zhang, M., Zhou, Z.Y. and Tan, J.W. (2010). Phenolic Compounds from *Merremia umbellata* subsp. *orientalis* and their allelopathic effects on *Arabidopsis* seed germination. *Molecules* **15**:8241-8250.